

## CACTIN pAb

<b>Catalog No.</b>	IDS0078
<b>Reactivity</b>	Human; Mouse; Rat
<b>Applications</b>	WB; ELISA
<b>Alternative Names</b>	Cactin; Renal carcinoma antigen NY-REN-24; CACTIN
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.07% sodium azide.
<b>Source</b>	Rabbit
<b>Dilution</b>	WB: 1:500-1:2000; ELISA: 1:10000-1:20000
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Concentration</b>	1 mg/ml
<b>Storage&amp;Stability</b>	Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.
<b>Subcellular Location</b>	-
<b>MW</b>	~ 95 kDa
<b>Background</b>	Involved in the regulation of innate immune response. Acts as negative regulator of Toll-like receptor, interferon-regulatory factor (IRF) and canonical NF-kappa-B signaling pathways. Contributes to the regulation of transcriptional activation of NF-kappa-B target genes in response to endogenous pro-inflammatory stimuli.
<b>Swiss-Prot</b>	Q8WUQ7

### Products Images:

