

Adrenocorticotropin(ACTH) (ABT-ACTH) mouse mAb Ready to use

Catalog No.	IML0065
Reactivity	Human
Applications	IHC-p
Gene Name	POMC
Protein Name	Pro-opiomelanocortin (POMC) (Corticotropin-lipotropin) [Cleaved into: NPP; Melanotropin gamma (Gamma-MSH); Potential peptide; Corticotropin (Adrenocorticotrophic hormone) (ACTH); Melanotropin alpha (Alpha-MSH); Corticotropin-like intermediary peptide (CLIP); Lipotropin beta (Beta-LPH); Lipotropin gamma (Gamma-LPH); Melanotropin beta (Beta-MSH); Beta-endorphin; Met-enkephalin]
Human Gene Id	5443
Swiss-Prot	P01189
Formulation	Liquid in PBS containing, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse:IgG1, Kappa
Dilution	Ready to use for IHC-p
Purification	The antibody was affinity-purified from mouse ascites by affinity-chromatography using specific immunogen.
Concentration	-
Storage&Stability	4°C: 1 years
Background	This gene encodes a preproprotein that undergoes extensive, tissue-specific, post-translational processing via cleavage by subtilisin-like enzymes known as prohormone convertases. There are eight potential cleavage sites within the preproprotein and, depending on tissue type and the available convertases, processing may yield as many as ten biologically active peptides involved in diverse cellular functions. The encoded protein is synthesized mainly in corticotroph cells of the anterior pituitary where four cleavage sites are used; adrenocorticotrophin, essential for normal steroidogenesis and the maintenance of normal adrenal weight, and lipotropin beta are the major end products. In other tissues, including the hypothalamus, placenta, and epithelium, all cleavage sites may be used, giving rise to peptides with roles in pain and energy homeostasis, melanocyte stimulation, and immune modulation.
Subcellular Location	Cytoplasmic
BiowMW	-

Products Images: