

PRODUCT DATA SHEET


TGFβ1 mouse Monoclonal Antibody(10E5)

Catalog No.	IMB0017
Reactivity	Human;Rat;Mouse
Applications	IHC-p;
Gene Name	TGFB1 TGFB
Protein Name	Transforming growth factor beta-1, TGF-β1, TGF b
Human Gene Id	7040
Swiss-Prot	P01137
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse
Dilution	IHC: 1:100-200
PurIF:ication	The antibody was affinity-purIF:ied from mouse ascites by affinity-
	chromatography using specIF:ic immunogen.
Concentration	1 mg/ml
Storage&Stability	-20°C/1 year
Background	This gene encodes a secreted ligand of the TGF-beta (transforming growth
	factor-beta) superfamily of proteins. Ligands of this family bind various TGF-
	beta receptors leading to recruitment and activation of SMAD family
	transcription factors that regulate gene expression. The encoded preproprotein
	is proteolytically processed to generate a latency-associated peptide (LAP)
	and a mature peptide, and is found in either a latent form composed of a mature
	peptide homodimer, a LAP homodimer, and a latent TGF-beta binding protein,
	or in an active form consisting solely of the mature peptide homodimer. The
	mature peptide may also form heterodimers with other TGFB family
	members. This encoded protein regulates cell prolIF:eration, dIF:ferentiation
	and growth, and can modulate expression and activation of other growth
	factors including interferon gamma and tumor necrosis factor alpha.
Subcellular Location.	[Latency-associated peptide]: Secreted, extracellular space, extracellular
	matrix. [Transforming growth factor beta-1]: Secreted.
BiowMW	-


Products Images:




PRODUCT DATA SHEET



Immunohistochemical analysis of paraffin-embedded Rat Brain Tissue using TGFβ1 Mouse mAb diluted at 1:200.



Immunohistochemical analysis of paraffin-embedded Human Breast Carcinoma Tissue using TGFβ1 Mouse mAb diluted at 1:200.



Immunohistochemical analysis of paraffin-embedded Mouse Lung Tissue using TGFβ1 Mouse mAb diluted at 1:200.