

## PRODUCT DATA SHEET


## P44/42 MAPK (ERK1/2) Monoclonal Antibody(6C10)

Catalog No.	IMB0046
Reactivity	Human;Rat;Mouse
Applications	WB; IHC-p
Gene Name	-
Protein Name	-
Human Gene Id	5594/5595
Swiss-Prot	P27361/P28482
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse
Dilution	WB: 1:1000-2000 IHC: 1:50-100
<b>PurIF:ication</b>	The antibody was affinity-purIF:ied from mouse ascites by affinity-
	chromatography using specIF:ic immunogen.
Concentration	1 mg/ml
Storage&Stability	-20°C/1 year
Background	The protein encoded by this gene is a member of the MAP kinase family. MAP
	kinases, also known as extracellular signal-regulated kinases (ERKs), act in a
	signaling cascade that regulates various cellular processes such as
	prolIF:eration, dIF:ferentiation, and cell cycle progression in response to a
	variety of extracellular signals. This kinase is activated by upstream kinases,
	resulting in its translocation to the nucleus where it phosphorylates nuclear
	targets. Alternatively spliced transcript variants encoding dIF:ferent protein
	isoforms have been described.
<b>Subcellular Location.</b>	Cytoplasm. Nucleus. Membrane, caveola. Cell junction, focal adhesion.
	Autophosphorylation at Thr-207 promotes nuclear localization
	(PubMed:19060905). PEA15-binding redirects the biological outcome of
	MAPK3 kinase-signaling by sequestering MAPK3 into the cytoplasm (By
	similarity).
BiowMW	-


## **Products Images:**



## PRODUCT DATA SHEET



Western blot analysis of 1)Mouse Brain Tissue, 2) Rat Brain Tissue with P44/42 MAPK(ERK1/2) Mouse mAb diluted at 1:2,000.



Immunohistochemical analysis of paraffin-embedded Human Breast Carcinoma using P44/42 MAPK (ERK1/2) Mouse mAb diluted at 1:200.