



## cIAP2 mouse mAb

<b>Catalog No.</b>	IMB0091
<b>Reactivity</b>	Human;Monkey
<b>Applications</b>	WB
<b>Gene Name</b>	birc3
<b>Protein Name</b>	-
<b>Human Gene Id</b>	330
<b>Swiss-Prot</b>	Q13489
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Monoclonal, Mouse
<b>Dilution</b>	WB: 1:1000
<b>Purification</b>	The antibody was affinity-purified from mouse ascites by affinity-chromatography using epitope-specific immunogen.
<b>Concentration</b>	1 mg/ml
<b>Storage&amp;Stability</b>	-20°C/1 year
<b>Background</b>	This gene encodes a member of the IAP family of proteins that inhibit apoptosis by binding to tumor necrosis factor receptor-associated factors TRAF1 and TRAF2, probably by interfering with activation of ICE-like proteases. The encoded protein inhibits apoptosis induced by serum deprivation but does not affect apoptosis resulting from exposure to menadione, a potent inducer of free radicals. It contains 3 baculovirus IAP repeats and a ring finger domain. Transcript variants encoding the same isoform have been identified. [provided by RefSeq, Aug 2011],
<b>Subcellular Location.</b>	Cytoplasm. Nucleus.
<b>BiowMW</b>	-

### Products Images:



Western blot detection of c-IAP2 in Ramos, COS7, Raji and Daudi cell lysates using c-IAP2 mouse mAb (1:1000 diluted). Predicted band size: 68KDa, Observed band size: 72KDa.



Western blot detection of cIAP2 in Ramos, COS7, Raji and HeLa cell lysates using cIAP2 antibody (1:1000 diluted).  
Exposure time: 5min. Predicted band size: 68KDa. Observed band size: 72KDa.