



## P44/42 MAPK (ERK1/2) Monoclonal Antibody(6B1)

<b>Catalog No.</b>	IMB0113
<b>Reactivity</b>	Human;Rat;Mouse
<b>Applications</b>	WB; IHC-p
<b>Gene Name</b>	-
<b>Protein Name</b>	-
<b>Human Gene Id</b>	5594/5595
<b>Swiss-Prot</b>	P27361/P28482
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Monoclonal, Mouse
<b>Dilution</b>	WB: 1:1000-2000 IHC: 1:50-100
<b>Purification</b>	The antibody was affinity-purified from mouse ascites by affinity-chromatography using specific immunogen.
<b>Concentration</b>	1 mg/ml
<b>Storage&amp;Stability</b>	-20°C/1 year
<b>Background</b>	The protein encoded by this gene is a member of the MAP kinase family. MAP kinases, also known as extracellular signal-regulated kinases (ERKs), act in a signaling cascade that regulates various cellular processes such as proliferation, differentiation, and cell cycle progression in response to a variety of extracellular signals. This kinase is activated by upstream kinases, resulting in its translocation to the nucleus where it phosphorylates nuclear targets. Alternatively spliced transcript variants encoding different protein isoforms have been described. [provided by RefSeq, Jul 2008],
<b>Subcellular Location.</b>	Cytoplasm. Nucleus. Membrane, caveola. Cell junction, focal adhesion. Autophosphorylation at Thr-207 promotes nuclear localization. PEA15-binding redirects the biological outcome of MAPK3 kinase-signaling by sequestering MAPK3 into the cytoplasm (By similarity).
<b>BiowMW</b>	-


### Products Images:



Western blot analysis of 1) Mouse Brain Tissue, 2) Rat Brain Tissue with P44/42 MAPK(ERK1/2) Mouse mAb diluted at 1:2,000.



Immunohistochemical analysis of paraffin-embedded Human Breast Carcinoma using P44/42 MAPK (ERK1/2) Mouse mAb diluted at 1:200.



The picture was kindly provided by our customer. Primary antibody was diluted at 1:2000. Loading control antibody was diluted at 1:20000