

PRODUCT DATA SHEET

TBP/TATA Binding Protein Monoclonal Antibody(1F6)


Catalog No.	IMB0143
Reactivity	Human;Rat;Mouse
Applications	IHC-p;
Gene Name	TBP
Protein Name	TATA-box-binding protein (TATA sequence-binding protein) (TATA-binding
	factor) (TATA-box factor) (Transcription initiation factor TFIID TBP subunit)
Human Gene Id	6908
Swiss-Prot	P20226
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse
Dilution	IHC: 1:50-200
PurIF:ication	The antibody was affinity-purIF:ied from mouse ascites by affinity-
	chromatography using specIF:ic immunogen.
Concentration	1 mg/ml
Storage&Stability	-20°C/1 year
Background	Initiation of transcription by RNA polymerase II requires the activities of more than 70 polypeptides. The protein that coordinates these activities is transcription factor IID (TFIID), which binds to the core promoter to position the polymerase properly, serves as the scaffold for assembly of the remainder of the transcription complex, and acts as a channel for regulatory signals. TFIID is composed of the TATA-binding protein (TBP) and a group of evolutionarily conserved proteins known as TBP-associated factors or TAFs. TAFs may participate in basal transcription, serve as coactivators, function in promoter recognition or modIF:y general transcription factors (GTFs) to facilitate complex assembly and transcription initiation. This gene encodes TBP, the TATA-binding protein. A distinctive feature of TBP is a long string of glutamines in the N-terminus. This region of the protein modulates the DNA bin
Subcellular Location.	Nucleus.

DIOWINI


Products Images:



PRODUCT DATA SHEET



Immunohistochemical analysis of paraffin-embedded Human Breast Carcinoma using TBP/TATA Binding Protein(mAb diluted at 1:200.



Immunohistochemical analysis of paraffin-embedded Mouse Kidney Tissue using TBP/TATA Binding ProteinMouse mAb diluted at 1:200.