


PRODUCT DATA SHEET

Histone H3 (Tri Methyl Lys9) Monoclonal Antibody(2C3)

Catalog No.	IMB0170
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	HIST1H3A/HIST1H3B/HIST1H3C/HIST1H3D/HIST1H3E/HIST1H3F/HIST1H3G/HIST1H3H
Protein Name	Histone H3.1/Histone H3.2/Histone H3.3
Human Gene Id	8350/8351/8352/8353/8354/8355/8356/8357/8358/8968
Swiss-Prot	P68431/Q71DI3/P84243
Formulation	PBS, pH 7.4, containing 0.5%BSA, 0.02% sodium azide as Preservative and 50% Glycerol.
Source	Monoclonal, Mouse
Dilution	WB: 1:1000-2000
PurIF:ication	The antibody was affinity-purIF:ied from mouse ascites by affinity-chromatography using specIF
Concentration	-
Storage&Stability	-20°C/1 year
Background	Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chrom of DNA wrapped around a nucleosome, an octamer composed of pairs of each of the four core he through the interaction of a linker histone, H1, with the DNA between the nucleosomes to for replication-dependent histone that is a member of the histone H3 family. Transcripts from this get This gene is found in the large histone gene cluster on chromosome 6p22-p21.3. [provided by Re
Subcellular	Nucleus. Chromosome.
Location.	
BiowMW	15273


Products Images:




Western blot analysis of lysates from 1) Hela, 2) Rat Testis tissue, 3) Raw264.7 cells, (Green) primary antibody was diluted at 1:1000, 4°over night, Dylight 800 secondary antibody(Immunoway:RS23910)was diluted at 1:10000, 37° 1hour. (Red) Actin β Polyclonal Antibody (Immunoway:YT0099) antibody was diluted at 1:5000 as loading control, 4° over night,Dylight 680 secondary antibody(Immunoway:RS23720)was diluted at 1:10000, 37° 1hour.



PRODUCT DATA SHEET



Western blot analysis of 1) Hela, 2) Rat Testis tissue, 3)
Raw264.7, diluted at 1:2000. cells nucleus extracted by Minute
TM Cytoplasmic and Nuclear Fractionation kit (SC003,Inventbiotech,MN,USA).



The picture was kindly provided by our customer