



## Cyclin E1 Polyclonal Antibody

<b>Catalog No.</b>	IPB0111
<b>Reactivity</b>	Human; Mouse; Rat
<b>Applications</b>	WB; IHC-p; IF/ICC; ELISA
<b>Dilution</b>	WB: 1:500-1:2000 IHC: 1:50-1:200 IF: 1:50-1:200 ELISA: 1:10000
<b>Gene Name</b>	CCNE1
<b>Protein Name</b>	G1/S-specific cyclin-E1
<b>Human Gene Id</b>	898
<b>Swiss-Prot</b>	P24864
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 05% BSA and 002% sodium azide
<b>Source</b>	Rabbit
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
<b>Concentration</b>	1 mg/ml
<b>Storage&amp;Stability</b>	-20°C/1 year
<b>Subcellular Location</b>	Nucleus
<b>MW</b>	47077
<b>Background</b>	The protein encoded by this gene belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance through the cell cycle Cyclins function as regulators of CDK kinases Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event This cyclin forms a complex with and functions as a regulatory subunit of CDK2, whose activity is required for cell cycle G1:S transition This protein accumulates at the G1-S phase boundary and is degraded as cells progress through S phase Overexpression of this gene has been observed in many tumors, which results in chromosome instability, and thus may contribute to tumorigenesis This protein was found to associate with, and be involved in, the phosphorylation of NPAT protein (nuclear protein mapped to the ATM locus), which participates in


### Products Images:




Qin, Zhenqi, et al. "Tsc1 deficiency impairs mammary development in mice by suppression of AKT, nuclear ER $\alpha$ , and cell-cycle-driving proteins." *Scientific reports* 6 (2016): 19587.




Western Blot analysis of various cells using Cyclin E1  
Polyclonal Antibody diluted at 1:1000



Western Blot analysis of customer's using Cyclin E1 Polyclonal  
Antibody. Antibody was diluted at 1:1000



Immunohistochemistry analysis of paraffin-embedded human  
colon carcinoma tissue, using Cyclin E1 Antibody. The picture  
on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from HeLa cells, treated with  
Paclitaxel 1uM 60', using Cyclin E1 Antibody. The lane on the  
right is blocked with the synthesized peptide.