

## ZMYM3 rabbit pAb

|                              |   |
|------------------------------|---|
| <b>Catalog No.</b>           | IPB14192  |
| <b>Reactivity</b>            | Human; Mouse;   |
| <b>Applications</b>          | WB  |
| <b>Dilution</b>              | WB: 1:500-2000  |
| <b>Gene Name</b>             | ZMYM3 DXS6673E KIAA0385 ZNF261  |
| <b>Protein Name</b>          | ZMYM3   |
| <b>Human Gene Id</b>         | 9203  |
| <b>Swiss-Prot</b>            | Q14202  |
| <b>Formulation</b>           | Liquid in PBS containing 50% glycerol, 05% BSA and 002% sodium azide  |
| <b>Source</b>                | Rabbit  |
| <b>Purification</b>          | The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen  |
| <b>Concentration</b>         | 1 mg/ml   |
| <b>Storage&amp;Stability</b> | -20°C/1 year  |
| <b>Subcellular Location</b>  | Nucleus   |
| <b>MW</b>                    | 150700  |
| <b>Background</b>            | This gene is located on the X chromosome and is subject to X inactivation It is highly conserved in vertebrates and most abundantly expressed in the brain The encoded protein is a component of histone deacetylase-containing multiprotein complexes that function through modifying chromatin structure to keep genes silent A chromosomal translocation (X;13) involving this gene is associated with X-linked mental retardation Several alternatively spliced transcript variants have been found for this gene |

### Products Images:

