

IDD 2050

PRODUCT DATA SHEET

IL-12B p40 Polyclonal Antibody

Catalog No.	IPB3850
Reactivity	Human; Mouse; Rat
Applications	WB; IHC; ELISA
Dilution	WB: 1:500-2000 IHC: 1:50-1:200 ELISA: 1:10000-20000
Gene Name	IL12B NKSF2
Protein Name	Interleukin-12 subunit beta (IL-12B) (Cytotoxic lymphocyte maturation factor
	40 kDa subunit) (CLMF p40) (IL-12 subunit p40) (NK cell stimulatory factor
	chain 2) (NKSF2)
Human Gene Id	3593
Swiss-Prot	P29460
Formulation	Liquid in PBS containing 50% glycerol, 05% BSA and 002% sodium azide
Source	Rabbit
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-
	chromatography using epitope-specific immunogen
Concentration	1 mg/ml
Storage&Stability	-20°C/1 year
Subcellular Location	Secreted
MW	-
Background	This gene encodes a subunit of interleukin 12, a cytokine that acts on T and

This gene encodes a subunit of interleukin 12, a cytokine that acts on T and natural killer cells, and has a broad array of biological activities Interleukin 12 is a disulfide-linked heterodimer composed of the 40 kD cytokine receptor like subunit encoded by this gene, and a 35 kD subunit encoded by IL12A This cytokine is expressed by activated macrophages that serve as an essential inducer of Th1 cells development This cytokine has been found to be important for sustaining a sufficient number of memory:effector Th1 cells to mediate long-term protection to an intracellular pathogen Overexpression of this gene was observed in the central nervous system of patients with multiple sclerosis (MS), suggesting a role of this cytokine in the pathogenesis of the disease The promoter polymorphism of this gene has been reported to be associated with the severity of atopic and non-atopic asthma in children


Products Images:




PRODUCT DATA SHEET



Western blot analysis of mouse-kidney lysate, antibody was diluted at 1000. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded humancolon, antibody was diluted at 1:200



Immunohistochemical analysis of paraffin-embedded humancolon, antibody was diluted at 1:200